

RECOMBINATION AND EXCITATION OF O_2^+ : CROSS SECTIONS AND PRODUCT YIELDS

R. Peverall^a, S. Rosen^b, J.R. Peterson^c, M. Larsson^b, A. Al-Khalili^b, L. Viktor^b, J. Semaniak^d, R. Bobbenkamp^c, A. Le Padellec^b, H. Danared^f, M. af Ugglas^f, and W.J. van der Zande^a

^aFOM Institute for Atomic and Molecular Physics, Kruislaan 407, 1098 Amsterdam, The Netherlands

^bDepartment of Physics, Stockholm University P.O. Box 6730, S-113 85 Stockholm, Sweden

^cMolecular Physics Laboratory, SRI International, Menlo Park, California 94025, USA

^dInstitute of Physics, Pedagogical University, ul. M. Konopnickiej 15, 25-430 Kielce, Poland

^eFaculty of Physics, University of Bielefeld, 33615 Bielefeld, Germany

^fManne Siegbahn Laboratory, Stockholm University S-104 05 Stockholm, Sweden

The cross sections for dissociative recombination and excitation of O_2^+ ($v=0$) have been measured in the ion storage ring CRYRING in Stockholm. The cross section gives a thermal rate coefficient of $2 \times 10^{-7} \text{ cm}^3 \text{ s}^{-1}$ at 300 K. An imaging technique was used to measure the $O(^1S)$ quantum yield, which is shown to have a surprisingly strong dependence on the electron energy.

Dissociative recombination (DR) and excitation of O_2^+ have been investigated using the ion storage ring CRYRING at the Manne Siegbahn Laboratory at Stockholm University. DR of O_2^+ plays an important role in atmospheric physics and chemistry, in particular in the F-region (above 140 km) where it is the source of the 630.0 nm red airglow and the 557.7 green airglow (arising from the $O(^1S) \rightarrow O(^1D)$ transition)

The green airglow has a long history of controversy¹ which is related to the problems to measure or control the vibrational population of O_2^+ ions either formed in the laboratory of the Earth's ionosphere, the problems to measure the quantum yield of $O(^1S)$ in the laboratory, and the problems of quantum calculations of DR of O_2^+ . In 1997 two important steps forward were taken. The $O(^1S)$ quantum yield for DR of O_2^+ populating a broad vibrational distribution was measured for an electron energy of nominally 0 eV using the ASTRID storage ring and an imaging technique². Despite the complication with using vibrationally hot O_2^+ , it was possible to deduce an $O(^1S)$ quantum yield of 0.05 for DR of O_2^+ ($v=0$)². A new mechanism was suggested³ and showed to quantitatively give a yield in quite good agreement with experiment².

Three experimental improvements were implemented in the present work. An ultracold electron beam allowed cross sections and quantum yields to be measured at an energy resolution of about 1 meV, a specially designed ion source supplied O_2^+ in its zeroth vibrational level, and an image intensifier was added to our three-dimensional imaging detector, which is similar in design to the one described in ref. 4 (in earlier version of our detector is described in ref. 5). The use of an image intensifier strongly reduced background events arising from collisions of O_2^+ in the rest gas.

Thus, for the first time one experiment

combines absolute cross section measurements with quantum yield determinations as a function of electron energy for vibrationally cold O_2^+ . Fig. 1 shows the DR cross section as a function of electron energy. Integration of the cross section in the usual way gives a thermal rate coefficient of $2 \times 10^{-7} \text{ cm}^3 \text{ s}^{-1}$ at 300 K which is in very good agreement with several earlier measurements¹. The $O(^1S)$ quantum yield shows a strong dependence on the electron energy.




Fig. 1 DR cross section for O_2^+

References

1. D.R. Bates, *Adv. At. Mol. Opt. Phys.* **34**, 427 (1994).
2. D. Kella, L. Vejby-Christensen, P.J. Johnson, H.B. Pedersen, and L.H. Andersen, *Science* **276**, 1530 (1997); *ibid.* **277** 167 (1997).
3. S.L. Guberman *Science* **278**, 1276 (1997).
4. Z. Amitay and D. Zajfman, *Rev. Sci. Instrum.* **68**, 1387 (1997).
5. S. Rosén, R. Peverall, J. ter Horst, G. Sundström, I. Semaniak, O. Sundqvist, M. Larsson, M. de Wilde, and W.J. van der Zande *Hyperfine Interact.* **115**, 201 (1998).