AU Le Padellec, A, Sheehan, C, Talbi, D, Mitchell, JBA

TI A merged-beam study of the dissociative recombination of HCO+

SO JOURNAL OF PHYSICS B-ATOMIC MOLECULAR AND OPTICAL PHYSICS

LA English

DT Article

ID STORAGE RING; IONS; H-3+; HEH+; COEFFICIENTS; IONIZATION; ELECTRONS; APPARATUS; STATE; CROSS

AB Cross sections for the dissociative recombination of HCO+ ions have been measured in the energy range from 0.01-1.0 eV. The large values measured for low-energy electrons are supported by an earlier extensive quantum mechanical study by Talbi and co-workers which is reviewed. A broad maximum has been found in the region from 0.3-0.8 eV and this is attributed to the opening up of a second product channel leading to CH + O.

C1 UNIV WESTERN ONTARIO, DEPT ANAT, DEPT PHYS & ASTRON,LONDON, ON N6A 3K7, CANADA. ENS, EQUIPE ASTROCHIM QUANT, LAB RADIOASTRON, F-75005 PARIS, FRANCE.

RP Le Padellec, A, UNIV RENNES 1, PALMS, CNRS, UMR 1203, F-35042 RENNES, FRANCE. NR 33

TC 8

PU IOP PUBLISHING LTD

PI BRISTOL

PA DIRAC HOUSE, TEMPLE BACK, BRISTOL, ENGLAND BS1 6BE

SN 0953-4075

J9 J PHYS-B-AT MOL OPT PHYS

JI J. Phys. B-At. Mol. Opt. Phys.

PD JAN 28

PY 1997

VL 30

IS 2

BP 319 EP 327

PG 9

SC Physics, Atomic, Molecular & Chemical; Optics

GA WJ025

UT ISI:A1997WJ02500023